



Study on biocontrol aspect of potential *Alcaligenes faecalis* against *Fusarium* sp., Concept and Approach

Purnima¹ and Pooja Singh²

¹Bacteriology and Natural Pesticide Laboratory, Department of Industrial Microbiology, Deen Dayal Upadhyaya Gorakhpur University, Gorakhpur- 273009, Uttar Pradesh, India

²Department of Botany, Deen Dayal Upadhyaya Gorakhpur University, Gorakhpur- 273009, U.P, India

Abstract

The world is dealing with vivid pollutions which lead to the degradation of resources and human health. In soil, the yield of crops is enhanced by the rampant use of chemical pesticides and fertilizers. The fungicides are applied on a large scale on almost all crops to control fungal infection which is the primary cause of crop productivity loss globally. The chemical fungicides are nevertheless causing great harm to soil quality and human health as many fungal pathogens are resistant to these chemical fungicides and all applied in bulk. There is a huge need for the substitution of such fungicides by natural components. The work is just an attempt in this aspect only to glorify *Alcaligenes faecalis* as a natural substitute for chemical fungicides as its bio-remediating aspect is explored earlier. The least explored association of *Alcaligenes faecalis* with *Fusarium* sp. which could be a futuristic fungicide is represented. *Fusarium* sp. (mainly *Fusarium oxysporum* and *Fusarium incarnatum*) are quite pathogenic causing great loss to various plants and crops across the globe. They are altogether responsible for wilt, leaf spot and root rot diseases. *Alcaligenes faecalis* as a potent biocontrol agent is introduced here as this bacterium has plant growth-promoting features too. So, *Alcaligenes faecalis* could become a natural, futuristic and biological fungicide and fertilizer. The large-scale bioinoculant production with its specific carrier molecules needs to be explored.

Keywords: *Alcaligenes faecalis*, *Fusarium oxysporum*, *Fusarium incarnatum*, antagonistic activity, biological fungicides, biocontrol agent, sustainable agriculture.

Alcaligenes faecalis: An emerging biocontrol agent

Alcaligenes faecalis was isolated from a wastewater bioprocessor in Texas, Houston, (USA) [1]. This genus is found in water, soil and in humans, generally non-pathogenic in nature. It is Gram-negative, aerobic, motile and rod-shaped which shows reactivation with Tryptone Soya broth (DSMZ medium 220a). It is mesophilic and shows optimum growth at 30-37°C. The genus is urease, gelatinase, arginine dihydrolase, beta-glucosidase negative, and cytochrome oxidase (enzyme activity) positive [2]. It belongs to Domain- Bacteria, Phylum- Pseudomonadota, Class- Betaproteobacteria, Order- Burkholderiales, Family- Alcaligenaceae, Genus- *Alcaligenes* and Species- *faecalis*.

Applicability of *Alcaligenes faecalis*

The bacteria has multitudinal aspects and cause bioremediation of phenanthrene, polyaromatic hydrocarbons, phenols, pesticides, and azo dyes. *Alcaligenes faecalis* also converts the most toxic form of arsenic, arsenite to a less fatal form, arsenate. It shows tolerance to heavy metals and has nanoparticle production, biocontrol, and nematicidal activity [3]. It also produces detergents, gum, and bioplastics [4]. This genus has high applicability still it remains under-represented and understudied at the whole genomic level [5].

Fusarium species: A virulent phytopathogen

Fusarium belongs to the Phylum Ascomycota, Family Nectriaceae, and has a broad host range.

120 various formae speciales showing specificity to the host species are determined. It causes damage and disease in cotton, date palm, banana, and other field crops including flowers and vegetables. The disease symptoms are distinguishable viz., vascular wilt, leaf epinasty, lower leaf yellowing, stunted growth, defoliation, and plant death eventually. The only measure to control this disease to a certain extent is the application of resistant varieties [6].

Fusarium oxysporum widely occurs among root region microbes. Mostly it is saprophytic but certain species cause root rot and wilting and are pathogenic. A new approach to biocontrol activity is the application of nonpathogenic *Fusarium* sp. which combats the pathogenic one [7].

Fusarium sp.: A specific study of diseases occurring in plants

The great loss in productivity of tomatoes is caused by wilt disease of *Fusarium oxysporum* f. sp. *lycopersici*. Still, no definite treatment for such a disease is reported. The pathogenic form of *Fusarium oxysporum* causes wilt disease classified under formae speciales depending on its host. The main reason for *Fusarium* wilt is its introduction in certain places rather than the local eruption of this pathogen. The soil population of the fungi is mainly ruled by asexual reproduction [8].

Musa spp. (Banana) is widely produced and consumed fruit. Panama disease (*Fusarium* wilt) causes a huge loss in the yield of bananas.

16 January 2024: Received | 03 March 2024: Revised | 19 March 2024: Accepted | 10 April 2024: Available Online

Citation: Purnima and Pooja Singh (2024). Study on biocontrol aspect of potential *Alcaligenes faecalis* against *Fusarium* sp., Concept and Approach. *Journal of Plant Biota*. DOI: <https://doi.org/10.51470/JPB.2024.3.1.34>

Purnima | purni9393ma@gmail.com

Copyright: © 2024 by the authors. The license of *Journal of Plant Biota*. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

The most effective measure to deal with this is again the use of resistant cultivars to a certain limit [9].

Vascular wilt of tomato is caused by *Fusarium oxysporum* f. sp. *lycopersici*. Symptoms include dark brown vascular tissues, wilting and apical discoloration leading to plant death. Wilt disease occurs due to mycotoxins, mycelial accumulation around the xylem, tylose production, and host defense inhibition [10]. (Table 1) [18]

Table 1: *Alcaligenes faecalis* S18 effect on *Fusarium* wilt disease severity of tomato cv. Rio Grande plants (60 days of inoculation with *Fusarium oxysporum* f. sp. *lycopersici*) than Control + (uninoculated with the pathogen and untreated)[18]

Bacterial isolate	Disease severity	Vascular browning extent (cm)	Plant height (cm)	Fresh plant weight (g)
Control +	3.4±0.2	8.2±0.6	15.8±0.2	4.4±0.4
<i>Alcaligenes faecalis</i> S18	0.2±0.1	0.4±0.2	30.6±2.2	9.63±0.9

Global chickpea productivity is adversely affected by *Fusarium* wilt. The soilborne pathogen (*Fusarium oxysporum* f. sp. *cicer*) has high survivability and great pathogenic variability [11].

Another globally affected crop of *Fusarium* wilt is cotton (*Gossypium* sp.). This disease was first reported in 1892 in the Alabama cotton field. The causal organism infects all species of cotton plants. *Fusarium oxysporum* f. sp. *vasinfectum* (FOV4) infects cotton in all stages of growth growing in neutral and acidic soil. The symptoms include wilting of the plant, marginal yellowing of leaf, stunting, damage and brown lining in the vascular region, and interveinal chlorosis followed by necrosis and death of the plant Fig 1 [12].



Fig 1: *Fusarium oxysporum* f. sp. *vasinfectum* infection symptoms in cotton [12]

- A. Wilting of cotton seedlings.
- B. Brown staining of vascular tissue.
- C. Streak of wilt infection in taproot.

Spinach, radish, lamb's lettuce, lettuce, wild rocket, and cultivated rocket production are adversely affected by the *Fusarium incarnatum equiseti* species complex. This leads to leaf spot disease in leafy vegetables [13]. *Fusarium incarnatum* results in crown rot disease in *Cucumis sativus*. White mycelia covered the young fruit leading to its withering. The disease symptom was similar to the fruit rot disease of *Botrytis cinerea* [14]. Postharvest disease, melon rot is caused by the *Fusarium incarnatum equiseti* species complex [15].

Alcaligenes faecalis as a biocontrol agent

Plant growth-promoting rhizobacteria are a group of beneficial rhizospheric bacteria that enhance plant growth and yield. The PGPR attribute of *Alcaligenes faecalis* for different crops has been least explored. It was reported by Jia R. et al (2022)[16], that *Alcaligenes faecalis* Juj 3 upon seed inoculation showed biocontrol activity against *Plasmidiophora brassicae* in cabbage and Chinese cabbage. It reduced clubroot disease in cabbage (51.4%) and Chinese cabbage (37.7%). It enhanced cabbage chlorophyll content (23.3%) and root length (49%). It also enhanced photosynthesis upon treatment in normal and *Plasmidiophora brassicae*-stressed environments. Antagonistic activity of *Alcaligenes faecalis* against *Fusarium oxysporum* was reported by Sayyed RZ and Chincholkar SB (2009)[17]. The antifungal activity occurred due to siderophore production by *Alcaligenes faecalis*. 75 µl siderophore-broth gave the best inhibition of *Fusarium oxysporum* [17].

Alcaligenes faecalis subsp. *faecalis* S18 gave the highest suppression of *Fusarium* wilt of tomato as observed by Abdallah RAB et al. (2016) [18]. This isolate also enhanced plant height, aerial part fresh weight, root length, and fresh weight. The antagonistic activity of the bacteria was due to chitinolytic, pectinolytic and hydrogen cyanide production Table 2 and Fig 2 [18].

Table 2: Antagonistic activity of *Alcaligenes faecalis* S18 against *Fusarium oxysporum* f. sp. *lycopersici* (FOL) [18]

Bacterial treatment	Colony diameter of FOL (cm)	Inhibition zone (mm)
Control (untreated)	3.71±0.08	0±0
<i>Alcaligenes faecalis</i> S18	2.15±0.05	12.37±0.1



Fig 2: Effects of *Alcaligenes faecalis* S18 on tomato plant growth parameters with and without FOL inoculation (60 days of harvesting). FOL: Stunted growth, FOL-S18: Good growth of plant [18]

Fusarium oxysporum f. sp. *cepae* growth was inhibited by 67.46% by consortia of *Bacillus mycoides* and *Alcaligenes faecalis* [19]. Biocontrol activity of *Alcaligenes faecalis* N1-4 against *Fusarium graminearum* and *Fusarium equiseti* was reported by Gong AD et al. (2019) [20].

Conclusions and Future Perspective

The work sheds light on the preambing to the antagonistic effect of *Alcaligenes faecalis*. Biocontrol aspect of *Alcaligenes faecalis* only against a *Fusarium* species viz, *Fusarium oxysporum* and *Fusarium incarnatum* are described as the most virulent species affecting global

crops and leading to great economic loss. Emphasis on biocontrol activity (PGPR attribute) of *Alcaligenes faecalis* is registered here as it is still under-explored. The most problematic content is no fungicides and other specific measures could combat this fungal pathogen.

There is a great demand for such bioinoculants and biological control agents in order to deal with the wilt and rot of *Fusarium* sp. The product should be cost-effective and eco-friendly and promote sustainable agriculture. The future of this study demands examining *Alcaligenes faecalis* inoculant against *Fusarium* sp. under field conditions on different crops and plants.

Conflict of Interest

The author has no conflict of interest in publishing this work.

References

1. Rehfuss M. and Urban J. 2006. *Alcaligenes faecalis* subsp. *phenolicus* DSM 16503. <https://www.dsmz.de/collection/catalogue/details/culture/DSM-16503>. Accessed on 23rd Dec, 2023.
2. Rehfuss M and Urban J. 2013. *Alcaligenes faecalis* subsp. *phenolicus* J is an aerobe, mesophilic, motile bacterium that was isolated from wastewater bioprocessor. <https://bacdiv.dsmz.de/strain/338>. Accessed on 23rd Dec, 2023.
3. Ju S., et al. *Alcaligenes faecalis* ZD02, a novel nematocidal bacterium with an extracellular serine protease virulence factor. *Appl. Environment Microbiology*. 2016. 82:2112-2120.
4. Tripathi AD., Srivastava SK., and Singh RP. Statistical optimization of physical process variables for bioplastics (PHB) production by *Alcaligenes* sp. *Biomass Bioenergy*. 2013. 55: 243-250. Doi- 10.1016/j.biombioe.2013.02.017.
5. Basharat Z., Yasmin A., He T., and Tong Y. Genome sequencing and analysis of *Alcaligenes faecalis* subsp. *phenolicus* MB207. *Scientific Reports*. 2018. 8: 3616. Doi- 10.1038/s41598-018-21919-4
6. Michielse CB., and Rep M. Pathogen profile update: *Fusarium oxysporum*. *Molecular Plant Pathology*. 2009. 10(3):311.
7. Fravel D., Olivain C., and Alabouvette C. *Fusarium oxysporum* and its biocontrol. *New Phytologist*. 2003. 157(3):493-502.
8. Widnyana K., et al. *Pseudomonas Alcaligenes*, potential antagonist against *Fusarium oxysporum* f. sp. *lycopersicum* the cause of *Fusarium* wilt disease on tomato. *Journal of biology, agriculture and healthcare*. 2013.3(7).
9. Ploetz RC. *Fusarium* wilt of banana. *Phytopathology*. 2015. 105(12): 1521-1521.
10. Srinivas C., et al. *Fusarium oxysporum* f. sp. *lycopersici* causal agent of vascular wilt disease of tomato. *Biology to diversity. A review. Saudi J. of biological sciences*. 2019. 26(7): 1315-1324.
11. Diaz RMJ., et al. *Fusarium* wilt of chickpeas: Biology, ecology and management. *Crop protection*. 2015. 73: 16-27.
12. Hu J., and Norton R. *Fusarium* wilt of Cotton. 2020. [Extension.arizona.edu/pubs/az1852-2020.pdf](https://extension.arizona.edu/pubs/az1852-2020.pdf). Accessed on 21st June, 2024.
13. Matic S., et al. Emerging leafy vegetable crop diseases caused by the *Fusarium incarnatum-equiseti* species complex. *Phytopathology Mediterr*. 2020. 59: 303-317.
14. Mao YS., et al. Occurrence of crown rot disease caused by *Fusarium incarnatum* on Cucumber (*Cucumis sativus*) in China. *Plant disease*. 2020. 104 (2):593.
15. Lima EN., et al. A novel lineage in the *Fusarium incarnatum-equiseti* species complex is one of the causal agents of *Fusarium* rot on melon fruits in Northeast Brazil. *Plant Pathology*. 2021. 70(1): 133-143.
16. Jia R., Chen J., Hu L., Liu X., Xiao K., and Wang Y. *Alcaligenes faecalis* Juj3 alleviates *Plasmodiophora brassicae* stress to cabbage via promoting growth and inducing resistance. *Front. Sustain. Food Syst*. 2022. 6. Doi- <https://doi.org/10.3389/fsufs.2022.942409>
17. Sayyed RZ., and Chincholkar. Siderophore-Producing *Alcaligenes faecalis* exhibited more biocontrol potential vis-à-vis chemical fungicide. *Current Microbiology*. 2009. 58(1):47-51.
18. Abdallah RAB., et al. Isolation of endophytic bacteria from *Withania somnifera* and assessment of their ability to suppress *Fusarium* wilt disease in tomato and to promote plant growth. *J. Plant Pathology Microbiology*. 2016. 7(5): 2-11.
19. Dinata GF., et al. In vitro evaluation of the effect of combined indigenous antagonistic bacteria against *Fusarium oxysporum*. *International J. of Agriculture System*. 2023. 11(1): 55-64.
20. Gong AD., et al. Inhibitory effect of volatiles emitted from *Alcaligenes faecalis* N1-4 on *Aspergillus flavus* and Aflatoxins in storage. *Frontiers in Microbiology*. 2019. 10: 1419.